

Acceleration Problems with Graphs


- 1. A bicyclist has an initial velocity of 2 m/s. Over 12 seconds, it speeds up to 8 m/s.
 - a. What was the accleration of the bicylclist?



- b. How many seconds did it take to reach a velocity of 6 m/s?
- c. Make a correct velcocity vs time graph for this motion.
- d. From the graph, how could you determine the acceleration?
- 2. Another bicylclist has an initial velocity of 16 m/s. They slow down to 4 m/s, in a time of 6 seconds.
 - a. What was the acceleration of the bicylclist?



- b. After only 3 seconds, what was their velocity?
- c. Make a correct position vs time graph for this motion.
- 3. The velocity vs time graph of something is shown to the right.
 - a. Describe the motion. (No calculations needed.)




t (s)

b. What is the acceleration during the first 8 seconds?

Acceleration Problems with Graphs

- c. What is the acceleration between 8 and 12 seconds?
- d. What is the acceleration for the last 12 seconds?
- e. What was the maximum speed of the object?
- f. When did the object have a velocity of 0? (Be careful!)
- 6. When is the object the farthest away from its original position? (You don't know how to calculate this, but just think about what the graph means.)
- 4. Starting from rest, a car has a constant acceleration of 3 mph/s for 6 seconds. It then has a constant speed for the next 3 seconds. Then it has a constant acceleration of -2 mph/s for the next 3 seconds.
 - a. After the first 6 seconds, what is the velocity of the car?



b. What is the acceleration of the car while it

has a constant speed?

- c. What was the velocity of the car after the last 3 seconds?
- d. Make an appropriate velocity vs time graph for this motion.

Answers:

1. a) 0.5 m/s² b) 8 s d) the slope of the velocity line is the acceleration

2. a) -2 m/s² b) 10 m/s 3. b) 1.25 m/s² c) 0 m/s² d) -0.417 m/s² e) 12.5 m/s

f) never g) t = 24 s 4. a) 18 mph b) 0 mph/s c) 12 mph