



Oscillation Problems III

1. A simple pendulum has a period of 1.5 seconds on the earth. If the same pendulum on another planet has a period of 3 seconds, what is the acceleration due to gravity on that other planet?

2. The angular position as a function of time for a simple pendulum is shown below.



- a. What is the length of the pendulum?
- b. What is the maximum linear speed of the mass at the end of the simple pendulum?
3. A simple pendulum of mass m and length L is hanging from a point "O." Directly underneath "O" is a pin "P" that is fixed in place. When the pendulum is released, the pin P becomes the new oscillation axis for that half of the motion. P is $L/2$ beneath O. What is the resulting period of small oscillations?




Oscillation Problems III

4. A physical pendulum is any body that is hung from a point (not its center of mass) and set oscillating back and forth. Calling the mass of the body " m " and the distance between the oscillation axis and the center of mass " r " and its moment of inertia about that axis " I ", what is the period of small oscillation for a physical pendulum?
5. What is the period of small oscillation for a thin rod of mass M and length L that is oscillating about one of its end points?

Oscillation Problems III

6. A thin rod of mass 400 grams and length 75 cm is suspended from one of its ends. At its other end is a small spring ($k = 125 \text{ N/m}$) attached horizontally to a wall. The system is in equilibrium when it is hanging vertically. What is the period of small oscillation?



Answers:

- 1) 2.5 m/s^2 2. a) 4.05 m b) 0.64 m/s 3) $\pi(1 + \sqrt{2})\sqrt{\frac{L}{2g}}$ 4) $2\pi\sqrt{\frac{I}{rmg}}$ 5) $2\pi\sqrt{\frac{2L}{3g}}$
- 6) 0.203 s